



Well Child Schedule*

Immunizations & Screening Tests

Newborn	Hepatitis B (if not given in hospital). Nirsevimab (RSV passive immunization) if desired.
2 weeks	None
1 month	None
2 months	Vaxelis (DTaP-IPV-Hep B-HIB), Prevnar 20 (pneumococcal), rotavirus.
4 months	Vaxelis (DTaP-IPV-Hep B-HIB), Prevnar 20 (pneumococcal), rotavirus.
6 months	Vaxelis (DTaP-IPV-Hep B-HIB), Prevnar 20 (pneumococcal), rotavirus.
9 months	Lead screening or test.
12 months	MMR, Varivax (chickenpox), Hepatitis A, Hemoglobin test +/- lead test.
15 months	DTaP, Hib, Prevnar 20 (pneumococcal).
18 months	Hepatitis A, lead screening or test
24 months	No vaccines if previous series completed; lead screening or test
3 years	No vaccines if previous series completed; lead screening or test
4 years	KG series: MMRV (MMR + Varivax), DTaP+IPV, lead screening or test
5 years	KG series (if not completed at 4 y/o), lead screening or test
6-10 yrs**	None. Recommend yearly check-up but no vaccines needed.
11 yrs**	MCV4 #1 (meningococcal), Tdap, HPV#1 (series of 2 if completed prior to age 15 years)
12 yrs	HPV #2
13-21 yrs**	Gardasil (HPV) if not completed
16 yrs	MCV #2, MenB #1 (meningococcal)
17 yrs	MenB #2
21 yrs	Tdap

We also recommend a **yearly Influenza vaccine** in the fall for all children age 6 months and older.

*We follow the Well Child and immunization schedules recommended by the American Academy of Pediatrics (AAP) and the Centers for Disease Control (CDC).

**A screening non-fasting lipid panel will be performed once between 9-11 years of age and again between 16-18 years of age. This is a recommendation of the AAP to screen for those at risk for early-onset heart disease.

Rev. 7/2024